

Survival of *Colletotrichum capsici* in decade-long cryopreserved Chilli (*Capsicum annuum*) seeds

USHA DEV, JAMEEL AKHTAR, REKHA CHAUDHURY¹, A. KANDAN, DINESH CHAND, JAGAT KUMAR, BALESHWAR SINGH, S.K. MALIK¹ AND P.C. AGARWAL

Division of Plant Quarantine, National Bureau of Plant Genetic Resources, New Delhi 110 012, India
devPusha26@rediffmail.com

Cryopreservation of orthodox seed germplasm from most common annual agronomic and horticultural species having desiccation and LN-tolerant has been carried out for more than 24 years at NBPGR [1, 2, 3]. The cryopreserved material is regularly monitored for seed viability and other parameters. Seed-health testing for pest-free conservation of germplasm at NBPGR was first initiated in 1998 for the material collected under National Agricultural Technology Project (NATP). In addition to quarantine of PGR under exchange, the Plant Quarantine Division of NBPGR also undertakes seed-health testing of germplasm conserved in the National Gene Bank (NGB) including cryogenebank material for pest-free conservation and further distribution. The present study reports observations recorded during seed health testing of cryopreserved chilli (*Capsicum annuum* L.) accessions.

A consignment comprising eight accessions of chilli seed was received by PQ Division in the month of September 2011 from the cryogenebank for seed-health testing prior to its release to the indenter. These accessions of chilli seeds were received at cryogenebank from NBPGR Regional Station, Hyderabad, India in 2001. The seed lots were desiccated to between 5 to 7% moisture contents before packing in 2 ml polypropylene cryovials and suspending at temperatures between -170⁰ to -180⁰C. The viability of fresh seed samples, as tested by Petri plate germination method [4], ranged from 84 to 92%. None of the

samples were given any physico-chemical treatments before storage. After 11 years of cryostorage the samples were thawed at room temperature and tested for viability and seed health.

During seed-health testing, all the chilli seed samples comprising 15-20 seeds, were first examined visually and then under stereo-binocular microscope for discolouration, deformation, malformation, fungal growth and fructifications, etc. Abnormal and unhealthy-looking seeds showing black spots were subjected to blotter test and examined on eighth day for presence of fungal pathogen(s) [5]. Fungi were identified on the basis of colony characters, fruiting bodies and conidia under stereo-binocular microscope and compound microscope. A single acervulus of the fungus on seed was picked up and transferred on potato dextrose agar (PDA) to isolate pure fungal cultures for ascertaining the identification [6].

During the dry seed examination of eight samples, only one sample, IC 214991 comprising 15 seeds, two seeds showed tiny black spots on seed surface. These spots in the blotter test developed into fungal colony showing dark acervuli (Fig.1a). Acervuli were sub-epidermal emerging by disrupting outer epidermal cell walls of host. Setae were dark brown, rigid, swollen at the base, slightly tapered to the paler acute apex, 1- to 5-septate, 250 x 6 µm (Fig.1b). Conidia were

¹Tissue Culture & Cryopreservation Unit, N.B.P.G.R., New Delhi 110 012, India

hyaline, falcate with acute apex and narrow truncate base, aseptate, uninucleate, $18-23 \times 3-5 \mu\text{m}$ (Fig.1c). Fungal colony on PDA was initially white which later became grey-dark brown. Mycelium formed light to dark grey cottony growth and acervuli with abundant dark setae. Based on the morphological characters of the acervuli, conidia and mycelial culture growing on the seed and PDA, the fungus was identified as *Colletotrichum capsici* (Syd.) Butler and Bisby [7].

The anthracnose fungus is reported to be seed-borne [8] and is known to survive in and on seed as acervulus and micro-sclerotia [9]. Survival of mycelia and stromata in colonized chilli seed had been reported by Manandhar *et al.* [10]. Mycelium is present in the outer and inner layers of the testa and in the endosperm which emerges to the surface after disrupting the seed coat [11]. Siddiqui *et al.* [12] reported the anthracnose fungus to be viable for over eight years in chilli seeds stored at 5°C . The pathogen can persist in soil, infected crop residues and weed [3]. The infected seeds give rise to weak seedlings which then become the primary source of inoculum in the field. During warm and wet season conidia from the acervuli get dispersed with wind or splashed by rain or irrigation water from diseased to healthy plants. All the samples were given fungicidal treatment with a mixture of mancozeb (0.2%) and bavistin (0.1%) as curative/prophylactic measure.

C. capsici which causes anthracnose disease in a wide range of hosts including cereals, legumes, vegetables, perennial crops and tree fruits, is one of the most important plant pathogens worldwide [3] and there has been a recent record of the fungus on *Jatropha curcus* [13]. Chilli is an important economic vegetable crop world wide, which gets severely infected by anthracnose causing yield losses up to 50% [14]. The disease is widely present in India and other chilli producing regions of the world [3]. Anthracnose disease complex of chilli is caused by more than one *Colletotrichum* species. Than *et al.* [15] have reported occurrence of different *Colletotrichum* species such as *C. acutatum* (Simmonds), *C. capsici*, *C. coccodes* (Wallr.) Hughes, *C. dematium* (Pers.) Grove, *C.*

gloeosporioides (Penz.) Penz. and Sacc. and *C. nigrum* (Wallr.) Hughes to be associated with anthracnose in chilli in various parts of the world. In India the disease is caused by *C. capsici* [16]. In addition to its wide distribution, there are reports of existence of variability in the pathogen [3]. Sharma *et al.* [17] reported existence of 15 pathotypes of *C. capsici*, from Himachal Pradesh in northern India based on the symptoms developed, whereas Montri *et al.* [18] reported three pathotypes in Thailand on the basis of quantitative infection on chilli. During last 3 decades, during processing 6398 samples of chilli germplasm imported from various countries, *C. capsici* was intercepted in seed samples from Nigeria, Taiwan, USA and Zambia [19].

The PGR are stored for utilization and posterity but full benefits of any storage system are realized only when the seeds intended for storage have high initial quality. Detection of *C. capsici* in the chilli seed conserved for more than ten years at temperature as low as -180°C shows that the fungus can survive even at this low temperature. These diseased seeds can act as a source of infection and can spread disease to new areas. Similar studies done on rice have proved that white tip nematode also survived in cryopreservation [20]. In view of India being the major chilli producing country, the disease has a great impact on chilli production, as the current commercial varieties are considered to be susceptible [21]. This further highlights the importance of seed-health testing in conserving disease-free material for quality assurance and minimizing the risk of spreading disease thus protecting Indian agriculture.

REFERENCES

1. MANDAL BB AND CHAUDHURY R (2004). Recent techniques for ex situ conservation of Plant Genetic Resources. In: Dhillon BS, Tyagi RK, Lal A and Saxena S (eds) *Plant Genetic Resources Management*. Indian Society of Plant Genetic Resources, New Delhi, Narosa Publishing House Pvt Ltd, New Delhi pp 194-204
2. CHAUDHURY R (2003). Principles of Cryopreservation. In: *In vitro Conservation and Cryopreservation of Tropical Fruit Species*. IPGRI Office for South Asia, New Delhi, India/ NBPGRI, New Delhi, India pp 125-32
3. MALIK SK, CHAUDHURY R, DHARIWAL OP

Received: March 2012

Revised: June 2012

Accepted: June 2012




Fig. 1. (a) Chilli seed showing acervulus of *Colletotrichum capsici*, (b) acervulus with setae, (c) conidia of *C. capsici*

- AND BHANDARI DC (2010). *Genetic Resources of Tropical Underutilized Fruits in India*. NBPGR, New Delhi, p 168
4. CHAUDHURY R, LAKHANPAUL S AND CHANDEL KPS (1989). Cryopreservation studies on plant germplasm. *Indian J Pl Genetic Resources* 2: 122-30
 5. AGARWAL PC, DEV USHA, SINGH B, RANI I, RAI D, JOSHI KD, CHAND D, MAURYA AK AND KHETARPAL RK (2004). Interception of seed-borne pathogens in introduced planting material. *Indian Journal of Microbiology* 44: 37-42
 6. GOH TK (1999). Single spore isolation using a hand-made glass needle. *Fungal Diversity* 2: 47-63
 7. ANONYMOUS (2007). *Crop Protection Compendium*. CAB International, Wallingford, UK
 8. RICHARDSON MJ (1990). *An Annotated List of Seed-borne Diseases*. ISTA, Zurich, Switzerland. p 387
 9. PERNEZNY K, ROBERTS PD, MURPHY JF AND GOLDBERG NP (2003). *Compendium of Pepper Diseases*. The American Phytopathological Society, St Paul, Minnesota, p 73
 10. MANANDHAR JB, HARTMAN GL AND WANG TC (1995). Anthracnose development on pepper fruits inoculated with *Colletotrichum gloeosporioides*. *Plant Disease* 79: 380-83
 11. MEON S AND NIK WZW (1988). Seed-borne infection and development of *Colletotrichum capsici* in naturally infected chili seed. *Pertanika*, 11(3): 341-44
 12. SIDDIQUI MR, MATHUR SB AND NEERGAARD P (1984). Longevity and pathogenicity of *Colletotrichum* spp. in seed stored at 5°C. *Seed Sci Technol* 11: 33-61
 13. TORRES-CALZADA C, TAPIA-TUSSELL R, NEXTICAPAN-GARCEZ A, MATIN-MEX R, QUIJANO-RAMAYO A, CORTES-VELAZQUEZ A, HIGUERA-CIAPARA I AND PEREZ-BRITO D (2011). First report of *Colletotrichum capsici* causing anthracnose in *Jatropha curcas* in Yucatan, Mexico. *New Disease Reports* 23: 6
 14. PAKDEEVARAPORN P, WASEE S, TAYLOR PWJ AND MONGKOLPORN O (2005). Inheritance of resistance to anthracnose caused by *Colletotrichum capsici* in *Capsicum*. *Plant Breeding* 124 (2): 206-08
 15. THAN PP, PRISHASTUTI H AND PHOULIVONG S (2008). Chilli anthracnose disease caused by *Colletotrichum* species. *Journal of Zhejiang University Science B* 9 (10): 774-78
 16. PAUL YS AND BEHL MK (1990). Some studies on bell pepper anthracnose caused by *Colletotrichum capsici* and its control. *Seed Res* 1: 656-59
 17. SHARMA PN, KAUR M, SHARMA OP, SHARMA P AND PATHANIA A (2005). Morphological, pathological and molecular variability in *Colletotrichum capsici*, the cause of fruit rot of chilies in the subtropical region of north-western India. *Phytopath* 153: 232-37
 18. MONTRI P, TAYLOR PWJ AND MONGKOLPORN O (2009). Pathotypes of *Colletotrichum capsici*, the causal agent of chili anthracnose in Thailand. *Plant Disease* 93 (1): 17-20
 19. AGARWAL PC, DEV USHA, SINGH B, RANI I, CHAND D AND KHETARPAL RK (2007). Seed-borne fungi identified from exotic pepper (*Capsicum* spp.) germplasm samples introduced during 1976-2005. *Plant Genetic Resources Newsletter* 149: 35-38
 20. RAJAN AND CHAUDHURY R (1996). Effect of storage temperature and moisture on the survival of the white tip nematode, *Aphelenchoides besseyi* and infested paddy, *Oryza sativa* seeds. *Ann Pl Protec Sci* 4: 16-21
 21. KAUR N, DHIMAN JS AND KHURANA DS (2011). Physiological and biochemical traits analysis of *Capsicum annuum* L germplasm for resistance to *Colletotrichum capsici*. *Cell and Plant Science* 23: 12-21.